arXiv:1507.06361v1 [math.GM] 23 Jul 2015

Characterizations of precompact sets in fuzzy star-shaped numbers space with L_p-metric

Huan Huang

Department of Mathematics, Jimei University, Xiamen 361021, China

---

1 Introduction

Compactness of fuzzy sets attracts many authors' attention. The readers may see Ref. [1–4,6,8,10] for some details. In this paper, we consider the characterizations of precompact sets in fuzzy star-shaped number space with L_p-metric.

2 Preliminaries

In this section, we introduce some basic notations and results. For details, we refer the readers to references [1, 9].

Let $\mathbb{N}$ be the set of all natural numbers, $\mathbb{R}^m$ be m-dimensional Euclidean space, $K(\mathbb{R}^m)$ be the set of all nonempty compact set in $\mathbb{R}^m$, and $C(\mathbb{R}^m)$ be the set of all nonempty closed set in $\mathbb{R}^m$. The well-known Hausdorff metric H on $C(\mathbb{R}^m)$ is defined by:

$$H(U,V) = \max\{H^*(U,V),\ H^*(V,U)\}$$

for arbitrary $U, V \in K(\mathbb{R}^m)$, where

$$H^*(U,V) = \sup_{u\in U} d\,(u,V) = \sup_{u\in U}\inf_{v\in V} d\,(u,v).$$

A set $K \in K(\mathbb{R}^m)$ is said to be star-shaped relative to a point $x \in K$ if for each $y \in K$, the line $\overline{xy}$ joining x to y is contained in K. The kernel ker K of K is the set of all points $x \in K$ such that $\overline{xy} \subset K$ for each $y \in K$. The symbol $K_S(\mathbb{R}^m)$ is used to denote all the star-shaped sets in $\mathbb{R}^m$.

Email address: `hhuangjy@126.com` .

Preprint submitted to Elsevier

$K_C(\mathbb{R}^m)$ is used to denote all the nonempty compact and convex sets in $\mathbb{R}^m$. Obviously, $K_C(\mathbb{R}^m) \subsetneqq K_S(\mathbb{R}^m)$.

We use $F(\mathbb{R}^m)$ to represent all fuzzy subsets on $\mathbb{R}^m$, i.e. functions from $\mathbb{R}^m$ to $[0,1]$. For $u \in F(\mathbb{R}^m)$, let $[u]_\alpha$ denote the α-cut of u, i.e.

$$[u]_\alpha = \begin{cases} \{x \in \mathbb{R}^m : u(x) \geq \alpha\}, \ \alpha \in (0,1], \\ \operatorname{supp} u = \overline{\{x \in \mathbb{R}^m : u(x) > 0\}}, \alpha = 0. \end{cases}$$

For $u \in F(\mathbb{R}^m)$, we suppose that
(i) u is normal: there exists at least one $x_0 \in \mathbb{R}^m$ with $u(x_0) = 1$;
(ii) u is upper semi-continuous;
(iii) u is fuzzy convex: $u(\lambda x + (1-\lambda)y) \geq \min\{u(x), u(y)\}$ for $x, y \in \mathbb{R}^m$ and $\lambda \in [0,1]$;
(iv) $[u]_\lambda$ is a star-shaped set for all $\lambda \in (0,1]$.
(v) $[u]_0$ is a compact set in $\mathbb{R}^m$.
(vi) $\left(\int_0^1 H([u]_\alpha, \{0\})^p \, d\alpha\right)^{1/p} < +\infty$, where $p \geq 1$ and 0 denotes the origin of $\mathbb{R}^m$.

- If u satisfies (i), (ii), (iii) and (v), then we say u is a fuzzy number. The set of all fuzzy numbers is denoted by E^m.
- If u satisfies (i), (ii), (iv) and (v), then we say u is a fuzzy star-shaped number. The set of all fuzzy star-shaped numbers is denoted by S^m.

Clearly, $E^m \subsetneqq S^m$. Diamond [1] introduced the d_p distance ($1 \leq p < \infty$) on S^m which are defined by

$$d_p\,(u, v) = \left(\int_0^1 H([u]_\alpha, [v]_\alpha)^p \, d\alpha\right)^{1/p}, \tag{1}$$

for all $u, v \in S^m$. They pointed out that d_p is a metric on S^m.

Wu and Zhao [10] considered the compactness criteria of E^m and S^m with d_p metric.

If we relax assumption (v) a little to obtain assumption (vi), then we obtain the following L_p–type noncompact fuzzy numbers or fuzzy star-shaped numbers.

Definition 2.1 If $u \in F(\mathbb{R}^m)$ satisfies (i), (ii), (iii) and (vi), then we say u is a L_p-type non-compact fuzzy number. The set of all such fuzzy numbers is denoted by $\widetilde{E}^{m,p}$. If there is no confusion, we also denote it by $\widetilde{E}^m$ for simplicity.

If $u \in F(\mathbb{R}^m)$ satisfies (i), (ii), (iv) and (vi) then we say u is a L_p-type non-compact fuzzy star-shaped number. The set of all such fuzzy star-shaped numbers is denoted by $\widetilde{S}^{m,p}$. If there is no confusion, we also denote it by $\widetilde{S}^m$ for writing convenience.

Clearly, $E^m \subsetneqq \widetilde{E}^{m,p}$, $S^m \subsetneqq \widetilde{S}^{m,p}$ and $\widetilde{E}^{m,p} \subsetneqq \widetilde{S}^{m,p}$.

It's easy to check that the d_p distance, $p \geq 1$, (see (1)) is also a metric on $\widetilde{S}^{m,p}$.

3 The characterization of precompact sets in $(\widetilde{S}^{m,p}, d_p)$

Definition 3.1 *[6] A set $U \subset \widetilde{S}^{m,p}$ is said to be uniformly p-mean bounded if there is a constant $M > 0$ such that $d_p(u, 0) \leq M$ for all $u \in U$.*

We can see that U is uniformly p-mean bounded is equivalent to U is a bounded set in $(\widetilde{S}^{m,p}, d_p)$.

Definition 3.2 *[1] Let $u \in \widetilde{S}^{m,p}$. If for given $\varepsilon > 0$, there is a $\delta(u, \varepsilon) > 0$ such that for all $0 \leq h < \delta$*

$$\left(\int_h^1 H([u]_\alpha, [u]_{\alpha-h})^p \, d\alpha\right)^{1/p} < \varepsilon,$$

where $1 \leq p < +\infty$, then we say u is p-mean left-continuous.

Suppose that U is a nonempty set in $\widetilde{S}^{m,p}$. If the above inequality holds uniformly for all $u \in U$, then we say U is p-mean equi-left-continuous.

Theorem 3.1 *U is a precompact set in $(\widetilde{S}^{m,p}, d_p)$ if and only if*
(i) U is uniformly p-mean bounded.
(ii) U is p-mean equi-left-continuous.

References

[1] P. Diamond, P. Kloeden, Metric Space of Fuzzy Sets−Theory and Application, World Scientific, Singapore, 1994.

[2] T. Fan, "On the compactness of fuzzy numbers with sendograph metric, Fuzzy Sets and Systems, vol. 143, no. 3,pp. 471C477, 2004

[3] Gabriele H. Greco, Sendograph metric and relatively compact sets of fuzzy sets, Fuzzy Sets and Systems 157 (2006) 286-291.

[4] H. Huang, C. Wu, A new characterization of compact sets in fuzzy number spaces, Abstract and Applied Analysis, Volume 2013, Article ID 820141, 6 pages, http://dx.doi.org/10.1155/2013/820141

[5] E.P. Klement, M.L. Puri, D.A. Ralescu, Limit theorems for fuzzy random variables, Proceedings of the Royal Society of London, Series A 407 (1986) 171-182.

[6] M. Ma, Some notes on the characterization of compact sets in (E^n, d_p), Fuzzy Sets and Systems 56 (1993) 297-301.

[7] C.V. Negoita, D.A. Ralescu, Application of fuzzy sets to system analysis. New York: Wiley, 1975.

[8] H. Román-Flores, The Compactness of $E(X)$, Appl. Math. Lett. Vol. 11, No. 2, pp. 13-17, 1998.

[9] C. Wu, M. Ma, The Basic of Fuzzy Analysis (in Chinese), National Defence Industry press, Beijing, 1991.

[10] C. Wu, Z. Zhao, Some notes on the characterization of compact sets of fuzzy sets with L_p metric, Fuzzy Sets and Systems 159 (2008) 2104-2115.